

THE STUDY OF PHYTOHORMONES AND EXPLANTS ON CALLUS INDUCTION AND REGENERATION OF SAINFOIN (*ONOBRYCHIS SATIVA*)

H.Garshasbi^{1,*}, M. Omid¹, S. Torabi¹ and D. Davodi.¹

Biotechnology Department, Agricultural College. Science and Research Branch. Islamic Azad Uni. , Tehran , Iran
Plant Breeding Department, Agricultural College , university of Tehran , karaj, Iran.

Research Center of Agricultural Biotechnology, Karaj, Iran.

*corresponding author's e.mail: garshasb2010@gmail.com

In this experiment, callus production and regeneration of sainfoin (*Onobrychis sativa* Var. Chadegan) were studied. The goal of this research was establishing a new method for micro proliferation of sainfoin. Following optimization of the growth regulator combinations in the Murashige and Skoog (MS) medium culture supplemented with combination of different 6-benzylaminopurine (BAP) and 2, 4- dichlorophenoxy acetic acid (2, 4-D) concentrations, a large number of shoots were propagated from four explants. The biggest callus was obtained on the MS medium containing 2, 4-D (6mg^{-1}) + BAP (1mg^{-1}) + Apical meristem explant with average 5.47 cm and the large number of shoots produced in the culture medium supplemented with BAP (4mg^{-1}) with average 1.6 plants per Petri dish (Bagheri and Safari 2004). Then, excised shoots rooted in half strength MS medium containing indole-3-butyric acid (IBA) (1mg^{-1}), transferred to perlite and grew into normal fertile plants (Sancak, 1998). This study showed that in callus induction of sainfoin meristem explant, BAP and 2-4, D together are important and in regeneration only BAP is effective.

Keywords: Callus production, regeneration, Sainfoin

INTRODUCTION

Sainfoins (*Onobrychis*) are Eurasian perennial herbs that have pale pink flowers and curved pods (Fig.1A). Its well-developed thick root system penetrates into the deeper layers of the soil; therefore, improves the organic matter of the soil and prevents soil erosion. It grows well in calcareous and chalky soils as well as in the soils with high water (Sancak, 1998). Sainfoin has needed no nitrogen fertilizer because there are many nitrogen stabilizing bacteria in its roots. It fattens livestock quicker than any other known forage. It has anthelmintic (anti-worm) properties and causes no bloat. Sainfoin is more tolerant to cold and drought; and more resistant to pests and diseases than alfalfa. Also, this plant is important in the honey industry. Breeding and biotechnology could be helpful in developing resistant varieties against different biotic and abiotic stresses with more flower size and yield. Regeneration via organogenesis or embryogenesis is the basis of tissue culture methods, and without regeneration it is impossible to produce transgenic plants. Efficient multiplication of this plant can offset lack of forage for growing livestock populations. Some of the varieties of sainfoin can grow in regions having unfavourable conditions for alfalfa's growth. The most ideal regeneration systems for genetic transformation are direct and repetitive production of somatic embryos or de novo shoot organogenesis originating from single cells in the epidermis (Rugkhla and Jones 1998). In a test, BAP at $22.2\text{ }\mu\text{M}$ supported the best combination of

shoot quality and number of shoots produced (Neves *et al.*, 2001). BAP and 2, 4- D together increased embryo induction while these couldn't influence them without each other (Nolan *et al.*, 2003). Auxin and cytokinin together increased callus, while BAP alone increased regeneration in some meristemic parts (Bagheri and Safari, 2004). H6 medium comprising MS salts with 2, 4-D (2mg^{-1}) and kinetin (0.5mg^{-1}) showed the greatest embryogenic efficiency (Lanas *et al.*, 2006). In somatic embryogenesis, somatic cells grow up as plants and this process is like zygotic embryogenesis (Gomes *et al.*, 2006). Whereas genetic clones created through shoot organogenesis require further manipulation for root initiation, in somatic embryogenesis, complete plants with a bipolar axis, vascular system and functional meristems are produced in a single step, (Bassuner *et al.*, 2007). In a tissue culture test, induction of nodular meristemoids from 86% of the leaf cultures was achieved on MS medium with 4.0 l M BA and 1.0 l M NAA. High levels (78-100%) of induction were also achieved on WPM with different densities of BA ($1.0\text{-}4.0\text{ l mg}^{-1}$) and IBA ($1.0\text{-}4.0\text{ l mg}^{-1}$). The highest conversion of meristemoids into shoots was only 22% for 4.0 l M BA and 1.0 l mg^{-1} NAA on MS initiation medium (Mayo *et al.* 2009). Recently, several reports have described shoot organogenesis was achieved through somatic embryo-like structures from different explants, mainly leaves and hypocotyls. In this research, a new method was developed for indirect organogenesis of sainfoin, two phytohormones

and 4 explants were applied, callus production and regeneration were examined.

MATERIALS AND METHODS

The present study was conducted in laboratory of Science and Research Branch, Islamic Azad University, Tehran, Iran during spring and summer 2006 to study and developed a new method for indirect organogenesis of sainfoin (*Onobrychis Sativa* Var. Chadegan).

Seed Sterilization and Germination: Seeds of sainfoin (*Onobrychis Sativa* Var. Chadegan) were surface sterilized in 100 % commercial bleach (containing 6% sodium hypochlorite) plus few drops of Tween 20 for 30 minutes with a continuous stirring, then rinsed three times with sterilized distilled water (Sancak1998) ; and inoculated in Petri dishes containing Agar solution (containing 7gr-1 Agar and 30g⁻¹ sucrose). Explants were obtained from sainfoin seedlings grown in plant growth chambers (Controlled Environments Ltd., Manitoba, Canada) at 25 ± 2°C under 16 hours photoperiod at a photosynthetic photon flux density of 100 µmol. m⁻². s⁻¹ provided by cool white fluorescent light (Mayo *et al*, 2009), After 7 to 10 days, seedlings grew 6 to 10 cm. Seedlings were cut in to 5×5 mm² segments and the explants comprised cotyledon leaf, apical meristem, stem and root.

Callus induction: MS medium (Murashige and Skoog, 1962) was supplemented with vitamins, sucrose (30 g-1),

myo-inositol (0.1 g-1), polyvinylpyrrolidone (3 g⁻¹) and plant growth regulators was adjusted to pH 5.6 before the addition of the gelling agent (8 g⁻¹ Agar Bacteriological, Agar No.1, Oxoid Ltd., Bastingstoke, England). The medium was then autoclaved at 121°C, 15 psi for 20 min. MS medium was poured in polystyrene Petri dishes (100 × 15 mm²) for callus induction and shoot regeneration. Grown Callus were transferred to Petri dishes after 37- 40 days. Callus divided into two pieces equal 5×5×5 mm³. The callus parts were transferred to new MS medium supplemented with the hormones for regeneration. Five callus of each were incubated in the photoperiod chamber for a month at 25±2°C, and then a number of plantlets were noted. The Callus production experiment was run as Factorial Randomized Complete Block Design with three factors and repetitions. The factors were explants (4 kinds), BAP and 2, 4-D. The densities of BAP and 2, 4-D were (0, 0.5, 1, 1.5 mg⁻¹) and (0, 2, 4, 6 mg-1), respectively. The Plant growth regulators were obtained from Sigma–Aldrich, St. Louis, USA.

Regeneration: The regeneration was the next stage of the experiment. callus divided into two pieces equal 5×5×5 mm³. The callus parts were transferred to new MSmedium supplemented with the hormones for regeneration. Five callus of each were incubated in the photoperiod chamber for a month at 25±2°C, and then a number of plantlets were noted (Fig.1.C).




Fig.1. A. Sainfoin's flower. B. A well developed callus of sainfoin (*Onobrychis sativa* Var. Chadegan) in MS medium containing two hormones, Bar = 1cm. C. Regeneration of sainfoin's calluses. D. Root production of plantlet in half-MS medium supplemented with IBA 91mgr.1⁻¹). E. Acclimatization of a sainfoin Plantlet (*Onobrychis sativa* Var. Chadegan) in pots covered with a plastic bag.

This part of experiment was conducted as Factorial Randomized Complete Block Design with 2 factors (BAP, 2, 4-D), in three repetition. The BAP densities were (0, 1.5, 4 mg^{-1}) and 2, 4-D concentrations were (0, 0.1, 3 mg^{-1}). Then, regenerated plantlets rooted with hormonal treatment (IBA 1 mg^{-1}) in half Ms medium for 4 weeks (Fig.1.D).

The plantlets were transferred to pots (covered with a plastic bag to prevent wilting for a few days) containing Pit most (very small particle that have no effect on plant and use only for root sustaining of plant), acclimatized in a greenhouse and then implanted to soil (Sancak, 1998). The data on callus production and regeneration were recorded to analysis of variance (ANOVA). The least significant difference test at 5 % Probable level was used for the mean separation.


RESULTS AND DISCUSION

The data emphasize the fact that sainfoin has a good capacity of *in vitro* callogenesis and regeneration (fig1.B). Many factors including the choice of growth regulators and choice of explants are responsible for successful callus production and regeneration. In the study of callus production about the leaf cotyledon explant, the greatest callus (5.411 cm) was produced in existence of two hormones with high ratio 2, 4-D to BAP, and the smallest callus produced in lack of 2, 4-D. These results confirm the previous findings of Bagheri and Safari (2004).

About the apical meristem, the greatest callus (5.477 cm) were produced in existence of two hormones with high ratio 2, 4-D to BAP, and the non calluses produced in the lack of two phytohormones or 2, 4-D. These results of this study were in agreement with the findings of Lanas *et al* (2006). Next, about the stem explant, the greatest callus (4.944 cm) grew in existence of two hormones with high ratio 2, 4-D to BAP, and the smallest callus (1.054 cm) produced in the lack of two growth plant regulators or BAP. These findings are in line with the finding of Mayo *et al*, (2009). Finally, in the root explant, maximum size of callus (5.208 cm) shown in existence of two hormones with high ratio 2, 4-D to BAP, and No callus was produced in the lack of BAP or 2, 4-D. These results confirm the previous finding of Bassuner *et al*. (2007) and Gomes *et al* (2006). The consideration on regeneration defined that BAP, 2, 4-D and BAP $\times$ 2, 4-D had sense in 1% probable levels. BAP had the most effect on regeneration (Graph 1, No7). Adding of BAP to 2, 4-D increased regeneration, (Graph 1). The BAP had positive and 2, 4-D negative effect on regeneration, and interaction effect between two hormones was negative (Graph 1). These results are in line with some scientists point of view, (Bagheri and Safari, 2004) and (Lanas *et al*, 2006).

The well developed shoots were transferred and cultured on half-strength MS medium supplemented with IBA (1 mg^{-1}).

More than 60% of shoots rooted on this medium within four weeks.



Graph 1 Effect of BAP $\times$ 2, 4-D interaction on regeneration of sainfoin levels BAP $\times$ 2,4-D (mg^{-1}) hormonal interaction
1: (0, 0), 2: (0, 0.1), 3: (0, 3), 4: (1.5, 0), 5: (1.5, 0.1), 6: (1.5, 3), 7: (4, 0), 8: (4, 0.1), 9: (4, 3).

CONCLUSION

Many factors, for successful callus production and regeneration, including the choice of growth regulators and choice of explants are important. The Increase of 2, 4-D (auxin) ratio to BAP (Cytokinin) raised callus production in each explant. The BAP and 2, 4-D had positive effect on the callus induction. The positive interaction effect was seen Between BAP and 2, 4-D. The Shoot meristemic explants produced the greatest calluses.

In the regeneration, it was determined that BAP had positive effect, 2, 4-D negative effect and there was the negative interaction effect between 2 hormones. This test and result are new because new species of sainfoin (*sativa*) was studied. Many factors including the choice of growth regulators and choice of explants are responsible for successful callus production and regeneration.

The results revealed that increase of 2, 4-D (auxin) (4, 6 mg^{-1}) ratio to BAP (Cytokinin) in each explant raised callus production of sainfoin. BAP and 2, 4-D had positive effect on the callus production. Between BAP and 2, 4-D was positive interaction effect. The Shoot meristemic explants produced the greatest callus.

In the regeneration, it was determined that BAP had positive effect, 2, 4-D had negative effect and between 2 hormones were negative interaction effect. 2, 4-D and BAP together had negative effect on regeneration.

ACKNOWLEDGEMENT

The authors appreciate the assistance of the laboratory personnel of Science and Research Branch, Islamic Azad University, Tehran.

REFERENCES

- Bagheri, A and M. safari. 2004. *In vitro* culture of higher plants. Ferdowsi Uni. Mashhad. p. 495.
- Bassuner, B., M. Lam, R. Lukowitz, W.E.C. Yeung. 2007. Auxin and root initiation in somatic embryos of Arabidopsis. *Plant Cell Rep.* 26:1-11.
- Gomes, F.L.A., F.F. Heredia, P.B. Silva, O. Faco, F.Campos. 2006. Somatic embryogenesis and plant regeneration in *Opuntia ficusindica* (L.) Mill. (Cactaceae). *Sci Hort.* 108:15-21.
- Lanas, I., P. Gallego, L. Martin, Fernandez, J. Alonso, A. J. Elena-Rosello, A. Blazquez, N.Villalobos and H. Guerra. 2006. In vitro culture of *Medicago arborea* L. anthers: Initial response. *Plant Growth Reg.* 49:49-60.
- Moyo, M., J. F. Finnie, J. Van Staden. 2009. In vitro morphogenesis of organogenic nodules derived from *Sclerocarya birrea* subsp. caffra leaf explants. *Plant Cell Tiss Org.* 98:273-280.
- Murashige, T., F. Skoog. 1962. A revised medium for rapid growth and bio assays with tobacco tissue cultures. *Physiol Plant* 15: 473-497.
- Neves, L. O., L. Tomaz and M. P. S. Fevreiro. 2001. Micropropagation of *Medicago truncatula* Gaertn. Cv. Jemalong and *Medicago truncatula* ssp. *Plant Cell Tiss Org.* 67:81-84.
- Omidi, M., B. Gerami and A. Moshari. 1992. Meristem culture in sainfoin. *Iranian J. of Agri. Sci.* 3: 49-57.
- Ozcan, S. C., S. Sevimay, M. Yddlz, C. Sancak and M. Ozgen. 1996. Prolific shoot regeneration from immature embryo explants of sainfoin (*Onobrychis viciifolia*). *Plant Cell Rep.* 16:200-203.
- Pupilli, F., F. Damiani, M. Pezzti and S. Arcioni. 1981. Plant regeneration from callus Protoplasts of *Onobrychis viciifolia* Scop (Sainfoin). *Plant Sci.* 63:87-94.
- Rugkhla, A., M.G.K. Jones. 1998. Somatic embryogenesis and plantlet formation in *Santalum album* and *S. spicatum*. *J. Exp. Bot.* 49:563-571.
- Salaj, J., B. Petrovska, B. Obert, A. Pret'ova. 2005. Histological study of embryo-like structures initiated from hypocotyl segments of flax (*Linum usitatissimum* L.). *Plant Cell Rep.* 24:590-595.
- Sancak, C. 1998. In vitro micropropagation of sainfoin (*O.vicifolia*). *Tubitak.* 23:133-136.
- Tamas, E and M. Savatti. 2006. In vitro micropropagation of sainfoin (*Onobrychis viclifolia* Scop.) *Agriculture.* 1:57-58.
- Tian, C., Y. Chen, X. Zhao and L. Zhao. 2008. Plant regeneration through protocorm-like bodies induced from rhizoids using leaf explants of *Rosa* spp. *Plant Cell Rep* 27:823-831.